

## NECK DISABILITY INDEX<sup>1</sup>

Section 1: To be completed by patient	AD	Non-Active Duty		
Name:	Age:	Date:		
Occupation:	Number of days of	f neck pain:	(this episode)	
Section 2: To be completed by patient				
This questionnaire has been designed to give your therapist information as to how your neck pain has affected your ability to manage in every day life. Please answer every question by placing a mark on the line that best describes your condition today. We realize you may feel that two of the statements may describe your condition, but <a href="please mark only the line">please mark only the line</a> <a href="which most closely describes your current condition">which most closely describes your current condition</a> .				
Pain Intensity				
I have no pain at the moment. The pain is very mild at the moment. The pain is moderate at the moment. The pain is fairly severe at the moment. The pain is very severe at the moment. The pain is the worst imaginable at the moment.	oment.			
Personal Care (Washing, Dressing, etc.)				
I do not have to change the way I wash an I do not normally change the way I wash o Washing and dressing increases my pain, Washing and dressing increases my pain, Because of my pain I am partially unable Because of my pain I am completely unab	or dress myself even to but I can do it without and I find it necessary to wash and dress with	hough it causes some pain. t changing my way of doin to change the way I do it. hout help.		
Lifting				
I can lift heavy weights without increased				
I can lift heavy weights but it causes incre Pain prevents me from lifting heavy weigh		it I can manage if they are o	conveniently	
positioned (ex. on a table, etc.).				
Pain prevents me from lifting heavy weights off of the floor, but I can manage light to medium weights				
if they are conveniently positioned.  I can lift only very light weights.				
I can not lift or carry anything at all.				
Reading				
I can read as much as I want to with no pa	in in my neck.			
I can read as much as I want to with slight				
I can read as much as I want with moderated I can't read as much as I want because of		neck.		
I can hardly read at all because of severe p	1			
I cannot read at all.				
Headache				
I have no headache at all.				
I have slight headaches which come infred				
I have moderate headaches which come in I have moderate headaches which come fi	1 2			
I have severe headaches which come frequency				
I have headaches almost all the time.		(Dan't forget to £11	out the best side)	
		(Don't forget to fill	out the back side)	



## NECK DISABILITY INDEX, p. 2

Section 2 (con't): To be completed by patient					
Concentration					
I can concentrate fully when I want to with no	difficulty.				
I can concentrate fully when I want to with slig	I can concentrate fully when I want to with slight difficulty.				
I have a fair degree of difficulty in concentrating	ing when I want to.				
I have a lot of difficulty in concentrating when					
I have a great deal of difficulty in concentrating	ng when I want to.				
I cannot concentrate at all.					
Work					
I can do as much as I want to.					
I can only do my usual work but no more.					
I can do most of my usual work, but no more.					
I cannot do my usual work.					
I can hardly do any work at all.					
I can't do any work at all.					
Driving Lead drive was a smith and a small main					
I can drive my car without any neck pain.  I can drive my car as long as I want with slight pain in my neck.					
I can drive my car as long as I want with moderate pain in my neck.					
I can't drive my car as long as I want because of moderate pain in my neck.					
	I can hardly drive at all because of severe pain in my neck.				
I can't drive my car at all.	,				
I have no trouble sleeping.  My sleep is slightly disturbed (less than 1 hour My sleep is mildly disturbed (1-2 hour sleep lo My sleep is moderately disturbed (2-3 hours sleep lo My sleep is greatly disturbed (3-5 hours sleep lo My sleep is completely disturbed (5-7 hours sleep lo My sleep is completely disturbed (5-7 hours sleep lo My sleep is completely disturbed (5-7 hours sleep lo My sleep is completely disturbed (5-7 hours sleep lo My sleep is completely disturbed (5-7 hours sleep lo My sleep is completely disturbed (5-7 hours sleep lo My sleep is completely disturbed (5-7 hours sleep lo My sleep is completely disturbed (5-7 hours sleep lo My sleep is completely disturbed (5-7 hours sleep lo My sleep is completely disturbed (5-7 hours sleep lo My sleep is greatly disturbed (3-5 hours sleep lo My sleep is greatly disturbed (3-5 hours sleep lo My sleep is greatly disturbed (3-5 hours sleep lo My sleep is greatly disturbed (5-7 ho	oss). leep loss). loss). leep loss). vities with no neck pain at all. vities with some pain in my neck. usual recreational activities because of pain in my neck. eational activities because of pain in my neck. ause of pain in my neck.				
Section 3: To be completed by physical therapist/pro					
SCORE:out of 50 (SEM 5, MDC 7)	Initial F/U weeks Discharge				
Number of treatment sessions:	Gender: Male Female				
Diagnosis/ICD-9 Code:					

<sup>&</sup>lt;sup>1</sup> Adapted from Vernon H, Mior S. The Neck Disability Indes: A Study of Reliability and Validitiy. Journal of Manipulative and Physiological Therapeutics 1991; 14(7): 409-415.



## NDI SCORING

The NDI is a modification of the Oswestry Low Back Pain Disability Index. The NDI can be scored as raw score (Vernon, 1991) or doubled, and expressed as a percent (Riddle, 1998). Each section is scored on a 0-5 scale, with the first statement being "0" (ie. No pain) and the last statement being "5" (ie. Worst imaginable pain). A higher score indicates more patient-rated disability. There is no statement in the original literature on how to handle missing data. To use the NDI for patient decisions, a clinically important change was calculated as 5 points, with a sensitivity of 0.78 and a specificity of 0.80 (Stratford, 1999).

Vernon H, Mior S. The neck disability index: a study of reliability and validity. J Manip Physiol Ther 1991; 14:407-415.

Riddle DL, Stratford PW. Use of Generic versus region specific functional status measures on patients with cervical spine disorders. Phys Ther 1998; 78:951-963.

Stratford PW. Riddle DL. Binkley JM. Spadoni G. Westaway MD. Padfield B. Using the neck disability index to make decisions concerning individual patients. Physiotherapy Canada, 107-112, 1999.